TOTONCHI
PLASTIC SURGERY

Care after Abdominoplasty (Tummy Tuck)

This page contains answers to frequently asked questions regarding your care after an abdominoplasty. This information will help you remember the instructions given to you after surgery.

Appearance:

- Your abdomen will feel tight and may appear mildly bruised and slightly swollen.
- Your original dressings and binder from surgery should remain on until your first shower, post-operative day 5. You will have two drains in place after surgery.
- We recommend that you wear an abdominal support garment for 6 weeks following your surgery, removing the garment only for bathing.
- Most patients find that a spandex or Lycra waist and tummy shaping garment are most comfortable to wear as a second stage garment, around 6 weeks post op.

Activity/Bathing/Sleeping:

- It will be difficult to stand up straight for several days. Assistance with ambulating may be helpful the first few days.
- Walk every day, but have someone assist you the first couple days.
- Depending on the type of work you do, you should be able to return to your job in 2-4 weeks or as soon as you are comfortable.
- Do not lift over 5 lbs or participate in heavy exercise for six weeks after the surgery.
- You generally should be able to drive about seven to ten days after surgery, if you feel comfortable. **You should never drive if you are still taking any prescription pain medication.**
- You can sponge bathe, but please **do not shower until your drains are removed.**
- Many patients find it easier to sleep in a recliner for a few days after the surgery. We encourage sleeping elevated for 1 week after surgery.

Drain/Incision Care:

- Your drains are held in place with stitches to your skin and should always be supported to avoid pulling on the drain. The easiest method is to safety pin the drain bulb to your abdominal binder or use the Velcro band provided.
- Monitor your drain output daily and record on the sheet given at your pre-operative appointment. Please bring the drain sheet to your post op appointments.
- The “How to care for your drain” pamphlet has more detailed information.
- Drains are generally removed within 7 to 10 days after your surgery.
- Clean your incisions with a Q-tip and warm water, then apply bacitracin to your incisions at least twice a day.

We are always available if you have any questions, please call 440-461-7999.

- After 1 week, please switch from bacitracin to Aquaphor or moisturizing lotion.
- Avoid sun exposure to the incision.
- Silicone scar gel can be applied to the scar at 5 weeks post-op. This will help aid the scar in healing.

Diet:

- We recommend a diet consisting of high protein and easily digestible foods (i.e. toast, rice, bananas, eggs, chicken, oatmeal, protein shakes, etc.)
- Do not drink alcohol for a couple weeks after your surgery or while taking narcotic pain medicines
- Limit your salt intake to help decrease swelling.
- Please refer to the “Medications and Foods to Avoid” List provided in your Pre-Surgical email sent

Medication:

- We will prescribe an antibiotic, pain medication, and nausea medication for you to take after your surgery. Follow the directions on the bottle for their use. The pain medication may make you feel drowsy
- A stool softener (Docusate) is recommended after surgery to ensure proper bowel regimen. If you have history of constipation, we advise starting the day before surgery and taking one tablet in the morning and evening.
- A week after surgery most patients will find that Tylenol will take care of the discomfort
- Do not use aspirin, ibuprofen (Motrin) or other NSAID medication for seven days after the surgery.

Post-Operative Appointments:

- **You will be seen 1 week after your surgery, then as indicated.**
- Additional post-op appointments will be scheduled as needed.

Recommended Supplies:

- Gauze, Q-tips, Bacitracin, and Aquaphor
- Extra Pillows for comfort
- Ice packs

Post-Operative Concerns: Urgent problems after abdominoplasty do not occur often. Please review the following complications that may require a change in your post-operative care:

- **Hematoma** (blood collecting under your incisions)

We are always available if you have any questions, please call 440-461-7999.

The warning signs are:
    - Severe pain that does not respond to medication
    - Significant swelling in your abdomen or localized burning
    - Excessive or growing bruising
- **Infection** is rare following abdominoplasty surgery. It is normal to have a small amount of drainage from the incisions and around your belly button for a few days. Signs of infection are:
    - Increased temperature
    - Increasing drainage from the incisions
    - Increasing redness around the incisions
- **Leg swelling** (with or without associated pain) may indicate a problem with the vein circulation in your leg(s). Although some swelling is to be expected due to your surgery, intravenous fluids given to you during the surgery, and your decreased level of activity, this swelling is normally mild, painless, and affects both legs evenly. If you have a large amount of leg swelling (either one or both legs), or if you experience pain in your legs, contact the office immediately.
- **Breathing problems** after abdominoplasty are rare but can be a serious complication. If you develop any chest and/or back pain or shortness of breath, you must contact my office or be seen in the nearest emergency medical facility.
- **Medication reactions** may occur with the drugs prescribed for you. If you develop a skin rash, itching, vomiting, or diarrhea, stop taking your medication and contact my office.

Stay Connected with Our Practice

For educational content, practice updates, and patient resources, follow Dr. Ali Totonchi on social media and visit our website:

Website: www.drtotonchi.com

Instagram: @totonchiplasticsurgery

Facebook: Dr. Ali Totonchi, MD

YouTube: Dr. Ali Totonchi

We are always available if you have any questions, please call 440-461-7999.